

# IMMUNOCHEMICAL LOCALIZATION OF HEPARANASE IN MOUSE AND HUMAN MELANOMAS

**Publication number:** JP5509403 (T)

**Publication date:** 1993-12-22

**Inventor(s):**

**Applicant(s):**

**Classification:**

**- international:** C07K7/08; C07K16/00; C07K16/30; C07K16/40; C12N9/24; G01N33/53; G01N33/573; G01N33/574; C07K7/00; C07K16/00; C07K16/18; C07K16/40; C12N9/24; G01N33/53; G01N33/573; G01N33/574; (IPC1-7): G01N33/53; C07K7/08; G01N33/573; C07K99/00

**- European:** C07K16/30L; C07K16/40; C12N9/24; G01N33/573; G01N33/574C12

**Application number:** JP19910512410T 19910530

**Priority number(s):** US19900530869 19900531


**Also published as:**

WO9119197 (A1)  
EP0532695 (A1)  
AU641269 (B2)  
AU8231791 (A)  
CA2083052 (A1)

Abstract not available for JP 5509403 (T)

Abstract of corresponding document: **WO 9119197 (A1)**

Antibodies directed against N-terminal heparanase peptide are produced. These antibodies are used for detection of heparan sulfate endoglycosidase in human and murine tumors. Localization and mounts of these heparanase antigens may ultimately be used to design appropriate therapeutic courses.



Data supplied from the esp@cenet database — Worldwide